

Act today to receive
a **\$100 rebate**
with your
second syringe of
JUVÉDERM®
injectable gel*



Follow the steps below to cash in on this exclusive offer from JUVÉDERM®


Step 1: Receive your JUVÉDERM® Ultra or JUVÉDERM® Ultra Plus treatment by **February 28, 2009**.

Step 2: Request the two syringe container lids from your treatment.

Step 3: Have your healthcare professional sign the form below to verify your treatment.

Step 4: Keep your receipt from your two-syringe JUVÉDERM® Ultra or JUVÉDERM® Ultra Plus injectable gel treatment.

Step 5: Mail your completed rebate form, the two syringe container lids, and the receipt by **March 31, 2009** to: **2009 Move to Two Rebate Offer, Delta Marketing Dynamics, One Clinton Square, 100 N. Salina St., Ste. 500, Syracuse, NY 13202.**



Syringe container lids, original receipt, and completed form must be submitted together.

*Second syringe must be administered on same treatment. Some restrictions may apply. Please see important treatment considerations on reverse side.

In the United States, JUVÉDERM® injectable gel is indicated for correction of moderate to severe facial wrinkles and folds (such as nasolabial folds).

ALLERGAN ©2009 Allergan, Inc.™ mark owned by Allergan, Inc. JUVÉDERM® mark owned by Allergan Industrie, SAS. THE SCIENCE OF REJUVENATION™ www.juvederm.com Re-order: MDOC-02323-04 900023



Yes, I want to cash in on the \$100 JUVÉDERM® rebate!

Name _____

Address _____

City _____ State _____ Zip code _____

Yes, I would like to receive future e-mail communications and special offers about JUVÉDERM® and other Allergan Medical Aesthetic products, programs, and services.

My e-mail address is: _____

To redeem this special, limited-time offer, please have your healthcare professional sign below:

Healthcare professional signature Date

Healthcare professional name (Please print)

Mail your completed rebate form, the two syringe container lids, and the receipt from your two-syringe JUVÉDERM® treatment to: 2009 Move to Two Rebate Offer, Delta Marketing Dynamics, One Clinton Square, 100 N. Salina St., Ste. 500, Syracuse, NY 13202.

Treatment must be received by 2/28/09 and the redemption forms must be postmarked by 3/31/09. Note: Rebate is valid on JUVÉDERM® Ultra and JUVÉDERM® Ultra Plus injectable gel only. Valid receipt and two syringe container lids must accompany form for redemption of this offer. This form may not be reproduced, duplicated, sold, or transferred in any way. This rebate cannot be combined with any other JUVÉDERM® offer. Only original rebate forms are valid and must accompany this request. Limit 1 reward per person. Maximum of 2 per household. Offer good only in the United States and Puerto Rico. Void where taxed or restricted. Please allow 4 to 5 weeks for delivery of your rebate check. If you have any questions about the rebate, call 1-800-492-4516.

A Brief Description of Relevant Indications for Use, Contraindications, Warnings, Precautions, and Adverse Events for JUVÉDERM® Injectable Gel

Indication: In the United States, JUVÉDERM® injectable gel is indicated for correction of moderate to severe facial wrinkles and folds (such as nasolabial folds).

Contraindications: JUVÉDERM® injectable gel should not be used in patients who have severe allergies marked by a history of anaphylaxis or history or presence of multiple severe allergies. JUVÉDERM® injectable gel should not be used in patients with a history of allergies to Gram-positive bacterial proteins.

Warnings: JUVÉDERM® injectable gel should not be injected into blood vessels. If there is an active inflammatory process or infection at specific injection sites, treatment should be deferred until the underlying process is controlled.

Precautions: The safety of JUVÉDERM® injectable gel for use during pregnancy, in breastfeeding females, or in patients under 18 years has not been established. The safety and effectiveness of JUVÉDERM® injectable gel for the treatment of areas other than facial wrinkles and folds (such as lips) have not been established in controlled clinical studies. Patients who are using substances that can prolong bleeding, such as aspirin or ibuprofen, as with any injection, may experience increased bruising or bleeding at injection site. Patients should inform their physician before treatment if they are using these types of substances. As with all skin-injection procedures, there is a risk of infection. JUVÉDERM®

injectable gel should be used with caution in patients on immunosuppressive therapy, or therapy used to decrease the body's immune response, as there may be an increased risk of infection. The safety of JUVÉDERM® injectable gel in patients with a history of excessive scarring (eg, hypertrophic scarring and keloid formations) and pigmentation disorders has not been studied. If laser treatment, chemical peel, or any other procedure based on active dermal response is considered after treatment with JUVÉDERM® injectable gel, or if JUVÉDERM® injectable gel is administered before the skin has healed completely after such a procedure, there is a possible risk of an inflammatory reaction at the treatment site.

Adverse events: The most commonly reported side effects are temporary injection-site redness, swelling, pain/tenderness, firmness, lumps/bumps, and bruising. Most side effects are mild or moderate in nature, and their duration is short lasting (7 days or less).

Important: For full safety information, please visit www.juvederm.com or call Allergan Product Support at 1-877-345-5372.

CAUTION: This device is restricted to sale by or on the order of a physician.
